

4032 FIRST AID KIT - 4032 first aid kit
Honeywell Safety Products USA, INC

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, [click here](#).

4032 First Aid Kit (Burn Relief Sp, EW, ASA, 1st aid Pack: triple, amm. Inh, BZK wipe-340001FP)

Eyewash

Active ingredient

Sterile Water 99%

Eyewash

Purpose

Eyewash

Eyewash

Uses

- for flushing the eye to remove loose foreign material, air pollutants or chlorinated water

Eyewash

Warnings

For external use only Obtain immediate medical treatment for all open wounds in or near eyes. To avoid contamination, do not touch tip of container to any surface. Do not reuse. Once opened, discard.

Do not use

- if solution changes color or becomes cloudy
- if you have open wounds in or near the eyes, get medical help right away.

Stop use and ask a doctor if

- you experience eye pain
- changes in vision
- continued redness or irritation of the eye
- condition worsens or persists

Keep out of reach of children

- If swallowed, get medical help or contact a Poison Control Center right away.

Eyewash

Directions

- remove contacts before using
- twist top to remove
- flush the affected area as needed
- control rate of flow by pressure on the bottle
- if necessary, continue flushing with emergency eyewash or shower

Eyewash

Inactive ingredients

sodium chloride, sodium phosphate dibasic, sodium phosphate monobasic

Eyewash

Questions

1-800-430-5490

Triple

Active ingredient (each gram contains)

Bacitracin zinc 400 units - Neomycin sulfate (5 mg equivalent to 3.5 mg Neomycin base) Polymyxin B sulfate 5000 units

Triple

Purpose

First aid antibiotic

First aid antibiotic

First aid antibiotic

Triple

Uses

first aid to help prevent infection in

- minor cuts
- scrapes
- burns

Triple

Warnings

For external use only

Allergy alert do not use if you are allergic to any of the ingredients

Do not use

- in the eyes
- over large areas of the body

Ask a doctor before use if you have

- a deep or puncture wounds
- animal bites
- serious burns

Stop use and ask a doctor if

- the condition persists or gets worse
- a rash or other allergic reaction develops
- you need to use longer than 1 week

Keep out of the reach of children

- If swallowed, get medical help or contact a Poison Control Center right away

Triple***Directions***

- clean the affected area
- apply a small amount of the product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily
- may be covered with a sterile bandage

Triple***Other information***

store at 15⁰ to 25⁰ C (59⁰ to 77⁰ F) tamper evident sealed packets - do not use if packet is torn or opened

Triple***Inactive ingredient***

petrolatum

Triple***Questions***

1-800-430-5490

Aspirin***Active ingredient (in each tablet)***

Aspirin 325 mg (NSAID)* *nonsteroidal anti-inflammatory drug

Aspirin***Purpose***

Pain reliever/fever reducer

Aspirin***Uses***

temporarily reduces fever and relieves minor aches and pains associated with:

- a cold
- headache
- toothache
- muscular aches
- backache
- minor pain of arthritis
- premenstrual and menstrual periods

Aspirin***Warnings***

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like

symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you are:

- age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for diabetes, gout or arthritis

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
 - pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present in the painful area
 - ringing in the ears or loss of hearing occurs
 - any new symptoms appear

If pregnant or breast-feeding,

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last three months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

- In case of overdose, get medical help or contact Poison Control Center right away.

Aspirin

Directions

- drink a full glass of water with each dose
- adults and children 12 years of age and older: take 1 or 2 tablets every 4 hours while symptoms last, not more than 12 tablets in 24 hours
- children under 12 years of age: consult a doctor

Aspirin

Other information

- store at room temperature 15° - 30°C (59° - 86°F)
- TAMPER EVIDENT PACKETS
- DO NOT USE IF OPEN OR TORN

Aspirin

Inactive ingredients

corn starch, croscarmellose sodium*, hypromellose*, microcrystalline cellulose*, mineral oil*, polyethylene glycol*, povidone, propylene glycol, silicon dioxide, stearic acid*, titanium dioxide*

*may contain these ingredients

Aspirin

Questions or Comments

1-800-430-5490

Ammonia

Active ingredient

Ammonia 15%

Ammonia

Purpose

Respiratory stimulant

Ammonia

Uses

- to prevent or treat fainting

Ammonia

Warnings

For external use only

Do not use

- if you have breathing problems such as asthma or emphysema

Stop use and ask a doctor if

- condition persists

Keep out of reach of children

- If swallowed get medical help or contact a Poison Control Center right away.

Ammonia

Directions

- hold inhalant away from face and crush ampoule between thumb and forefinger at position indicated on sleeve.
- hold near nostrils for inhalation of volatile vapor

Ammonia

Other information

- store at room temperature away from light

Ammonia

Inactive ingredient

alcohol USP, FD&C red #40, lavender oil, lemon oil fcc, nutmeg oil, purified water

Ammonia

Questions or Comments?

1-800-430-5490

BZK Wipes

Active ingredient

Benzalkonium chloride 0.13% w/v

BZK Wipe

Purpose

First aid antiseptic

BZK Wipes

Uses

- Antiseptic cleansing of face, hands, and body without soap and water

BZK Wipes

Warnings

For external use only

Do not use

- in the eyes or over large areas of the body
- on mucous membranes
- on irritated skin
- in case of deep puncture wounds, animal bites or serious burns, consult a doctor
- longer than 1 week unless directed by a doctor

Stop use and ask a doctor if

- if irritation, redness or other symptoms develop
- the condition persists or gets worse

Keep out of reach of children

- If swallowed, get medical help or contact a Poison Control Center right away.

BZK Wipe***Directions***

- tear open packet and use as a washcloth

BzK Wipe***Other information***

- store at room temperature 15⁰ to 30⁰ C (59⁰ - 86⁰ F)
- do not reuse towelette

BZK Wipe***Inactive ingredient***

water

BZK Wipe***Questions***

1-800-430-5490

Burn Relief***Active ingredient***

Benzethonium chloride 0.2% w/w

Benzocaine 10% w/w

Menthol 0.33% w/w

Burn Relief Spray***Purpose***

Topical antiseptic

Topical anesthetic

Topical anesthetic

Burn Relief Spray***Uses***

for the temporary relief of pain and itching and helps protect against infection in:

- minor cuts and scrapes
- burns
- sunburn
- insect bites
- minor skin irritations

Burn Relief Spray

Warnings

For external use only

Flammable

keep away from fire or flame

contents under pressure

do not puncture or incinerate container

do not expose to temperatures above 120 0 F

Do not use

- in or near the eyes or other mucous membranes
- in case of serious burns
- in case of deep or puncture wounds
- for prolonged period of time
- on large portion of the body

Stop use and ask a doctor if:

- condition worsens or symptoms persist for more than 7 days
- condition clears up and recurs within a few days
- redness, swelling, or irritation occurs

Keep out of the reach of children

If swallowed, get medical help or contact a Poison Control Center right away.

Burn Relief Spray

Directions

- clean the affected area
- shake can well before using
- hold 4 - 6 inches from surface and spray area until wet
- may be covered with a sterile bandage, if bandaged let dry first
- for adult institutional use only
- not intended for use on children

Burn Relief Spray

Other information

- avoid inhaling
- use only as directed
- intentional misuse by deliberately concentrating or inhaling the contents may be harmful or fatal

Burn Relief Spray

Inactive ingredients

dipropylene glycol, isobutane, n-butane, propane

4032

340001FP Kit Contents

1 3/4X3 PLAS SING 50/BOX
1 GAUZE BANDAGE, 4" X 6 YD
1 ADH TAPE W/P 1/2"X 2 1/2 YD
1 O/H PUMP BURN RELIEF 2 OZ ID G
1 GAUZE CLEAN-WRAP BDGE N/S 2"
1 1 OZ, BUFF EYEWASH
1 SCISSOR BDGE 4" RED PLS HDL
1 KIT TWEEZER 3 1/2" SLANTED
1 F. A. INST CHART SM (INDIVIDUAL LBL)
LBL STOCK 6-3/8"X4"
1 LBL STOCK 3"x1-7/8"
2 BAYER 12 PACK PER ZIP BAG
1 1ST AID PACK FOR 0001 0002 110
1 KIT, PP 10 UNIT FA

Eyewash

Principal Display Panel

Honeywell

TAMPER-EVIDENT CAP.
TAPA CON SELLO DE SEGURIDAD.
BOUCHON INDICATEUR D'INFRACTION.

eyesaline®

LAVAOJOS
EYESALINE

Solución
Isotónica Estéril

EYESALINE
EYEWASH

Sterile
Isotonic Solution

LAVAGE
OCULAIRE
EYESALINE

La Solution
Isotonique Stérile

16 fl. oz. (473 mL)

NPN: 80057528



Drug Facts (for USA only)

Active ingredient Sterile water 99% **Purpose** Eyewash
Uses for flushing the eye to remove loose foreign material, air pollutants, or chlorinated water.
Warnings For external use only - Obtain immediate medical treatment for all open wounds in or near the eyes. To avoid contamination, do not touch tip of container to any surface. Do not reuse. Once opened, discard.
Do not use if solution changes color or becomes cloudy
if you have open wounds in or near the eyes, get medical help right away
Stop use and consult a doctor if:
• you experience eye pain • changes in vision
• continued redness or irritation of the eye
• condition worsens or persists
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.
Directions
• remove contacts before using • twist top to remove
• flush the affected area as needed
• control rate of flow by pressure on the bottle
• if necessary, continue flushing with emergency eyewash or shower
Inactive ingredients sodium chloride, sodium phosphate dibasic, sodium phosphate monobasic
Questions? Call 1-800-430-5490
Honeywell Safety Products USA, Inc. Smithfield, RI. 02917

LABEL #32-0045/0 Rev. J REORDER / NUEVO PEDIDO / REAPPROVISIONNEMENT #32-000454-0000

space for lot code and supplier part number

PEEL / PELAR / PELEL


Datos de medicamento (Para EE.UU. solamente)

Ingrediente Activo Agua estéril 99% **Propósito** Lavaojos
Usos para el lavado de ojo para quitar las partículas sueltas y extrañas, los contaminantes aeros, o agua de cloruro
Advertencias Para el uso externo sólo - Obtenga tratamiento médico inmediato para todas las heridas abiertas en o cerca de los ojos. Para evitar la contaminación, no toque la punta del envase con ninguna superficie. No vuelva a usar. Vez abierto, descarte.
No se use si la solución se enturbia o cambia de color
si tiene heridas abiertas en o cerca del ojo, obtenga ayuda medica de inmediato
Deje de usar y consulte a un médico si:
• experimenta dolor de ojo • cambio de visión
• rojez continuo o irritación del ojo
• la condición empeora o persiste
Manténgase fuera del alcance de los niños. En caso de ingestión accidental, obtenga atención médica o llame a un Centro de control de envenenamiento inmediatamente.
Instrucciones
• quite los lentes de contacto antes de usar la solución
• tuerza la tapa para quitar
• enjuague el área afectada según se necesite
• controle el chorro haciendo presión el la botella
• si es necesario, sigue enjuagado con un lavaojos o ducha de emergencia
Ingredientes inactivos cloruro de sodio, fosfato de sodio dibásico, fosfato de sodio monobásico
¿Preguntas? Llame al 1-800-430-5490
Honeywell Safety Products USA, Inc. Smithfield, RI. 02917

Information

Usages Pour le rinçage des yeux afin d'enlever un corps étranger, des polluants atmosphériques ou de l'eau chlorée.
Advertissements Pour usage externe seulement - Obtenir immédiatement des soins médicaux pour toutes les plaies ouvertes dans ou près des yeux. Pour éviter toute contamination, ne pas toucher la pointe du récipient à n'importe quelle surface. Ne pas réutiliser. Une fois ouvert, jetez-le.
Ne pas utiliser si la solution a changé de couleur ou si elle est brouillée
si vous avez des plaies ouvertes aux yeux ou à proximité, consultez immédiatement un médecin
Cesser d'utiliser la solution et consulter un médecin si vous ressentez une douleur oculaire si votre vision change
rougeur ou irritation persistante des yeux
condition empire ou persiste
Garder hors de la portée des enfants. En cas d'ingestion, communiquer immédiatement avec un médecin ou avec un centre antipoison.
Mode d'emploi
• enlever les verres de contact avant l'utilisation • dévisser le bouchon pour l'enlever • rincer la zone touchée selon les besoins • ajuster le débit d'écoulement de la solution en augmentant ou en réduisant la pression exercée sur le contenant
• si nécessaire, continuer de rincer avec une solution de rinçage oculaire d'urgence ou une douche
Ingédients eau stérile, chlorure de sodium, phosphate dibasique de sodium, phosphate monobasique de sodium
Des questions? Faites le 1-800-430-5490
Honeywell Safety Products USA, Inc. Smithfield, RI. 02917

Triple
Principal Display Panel



Honeywell

**TRIPLE
ANTIBIOTIC
OINTMENT**

Bacitracin zinc, Neomycin sulfate,
Polymyxin B sulfate
First Aid Antibiotic

**Triple
Antibiotic
Ointment**

To Help Prevent Infection in
Minor Cuts, Scrapes and Burns

144 Packets
Net wt. per packet 0.5g (0.02 oz)

Honeywell

Honeywell

Honeywell

Honeywell

Drug Facts

Active Ingredients (in each gram)	Purpose
Bacitracin zinc 400 units	First aid antibiotic
Neomycin sulfate 5 mg (equivalent to 3.5 mg neomycin base)	First aid antibiotic
Polymyxin B sulfate 5000 units	First aid antibiotic

Uses

- first aid to help prevent infection in:
- minor cuts • scrapes • burns

Warnings

For external use only.

Allergy alert • do not use if you are allergic to any of the ingredients.

Do not use

- in the eyes
- over large areas of the body

Ask a doctor before use if you have

- deep or puncture wounds
- animal bites
- serious burns

Stop use and ask a doctor if

- condition persists or gets worse
- rash or other allergic reaction develops
- you need to use longer than 1 week

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- apply to affected area
- apply a small amount of product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily
- may be covered with a sterile bandage

Other information

- keep out of reach of children (39 to 79°F)
- tamper evident sealed packets
- do not use if packet is broken or opened

Inactive ingredient petrolatum

Questions? 1-800-438-5689

**TRIPLE
ANTIBIOTIC
OINTMENT**

Bacitracin zinc, Neomycin sulfate,
Polymyxin B sulfate
First Aid Antibiotic

**Triple
Antibiotic
Ointment**

To Help Prevent Infection in
Minor Cuts, Scrapes and Burns

Bacitracin zinc, Neomycin sulfate,
Polymyxin B sulfate

144 Packets
Net wt. per packet 0.5g (0.02 oz)

Distributed by
**Honeywell Safety
Products USA, Inc.**
Smithfield, RI 02917

Re-order #231209G

Caution:

This package is for industrial use only.
Not for retail sales.

Keep this and all drugs out of the reach of children.
In case of ingestion, seek professional assistance or
contact a poison control center immediately.

**TRIPLE
ANTIBIOTIC
OINTMENT**

Bacitracin zinc, Neomycin sulfate,
Polymyxin B sulfate
First Aid Antibiotic

**Triple
Antibiotic
Ointment**


To Help Prevent Infection in
Minor Cuts, Scrapes and Burns

Bacitracin zinc, Neomycin sulfate,
Polymyxin B sulfate

144 Packets
Net wt. per packet 0.5g (0.02 oz)

771209G Rev A

**Aspirin
Principal Display Panel**



7722-01 Rev C

Drug Facts (continued)
Active ingredient (in each tablet): Aspirin 325mg (NSAID)
Purpose: Pain reliever / fever reducer
Nonsteroidal anti-inflammatory drug
Uses: temporarily reduces fever and relieves minor aches and pains associated with:
 - cold - headache - toothache - menstrual aches - backache - minor pain of arthritis
Warnings:
 - aspirin allergy: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, changes in behavior with nausea and vomiting occur. Consult a doctor because these symptoms could be an early sign of Reye's Syndrome, a rare but serious illness.
Directions: See accompanying package inserts for directions.
Other information: Aspirin 325mg (NSAID)
Contains: Pain reliever / fever reducer

PACKAGE NOT CHILD-RESISTANT

25 Packets (2 Tablets per Packet)
 Total quantity 50 Tablets

ASPIRIN 325 MG
(NSAID)
Pain Reliever / Fever Reducer

Honeywell



Distributed by
Honeywell Safety Products, USA, Inc.
 Smithfield, RI 02917

35222AA

ASPIRIN 325 MG
 25 Packets (2 Tablets per Packet)
 Total quantity 50 Tablets

35222AA

Drug Facts (continued)
Allergy alert: Aspirin may cause a severe allergic reaction which may include:
 • hives • facial swelling • asthma (wheezing) • shock
Stomach bleeding warning: This product contains an NSAID which may cause severe stomach bleeding. The chance is higher if you:
 • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed
Do not use if you are allergic to aspirin or any other pain reliever/fever reducer.
Ask a doctor before use if: - stomach bleeding warning applies to you - you have a history of stomach problems such as heartburn - you have high blood pressure, heart disease, liver cirrhosis, or kidney disease - you are taking a diuretic - you have asthma
Ask a doctor or pharmacist before use if you are: taking a prescription drug for diabetes, gout or arthritis
Stop use and ask a doctor if: - you experience any of the following signs of stomach bleeding:
 • feel faint • vomit blood • have bloody or black stools • have stomach pain that does not get better
 • pain gets worse or lasts more than 10 days • fever gets worse or lasts more than 3 days
 • redness or swelling is present in the painful area • ringing in the ears or loss of hearing occurs
 • any new symptoms appear
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center immediately.



Drug Facts (continued)
Directions - drink a full glass of water with each dose
 • adults and children 12 years of age and older: take 1 - 2 tablets every 4 hours while symptoms last, not more than 12 tablets in 24 hours
 • children under 12 years of age: consult a doctor
Other information - Store at room temperature 15° - 30° C (59° - 86° F)
 • **TRIPLEX EVIDENCE PACKETS - DO NOT USE IF OPEN OR TORN**
Inactive ingredients - corn starch, croscarmellose sodium, hypromellose, microcrystalline cellulose, mineral oil, polyethylene glycol, povidone, propylene glycol, silicon dioxide, stearic acid, titanium dioxide
 *may contain these ingredients
Questions or comments? 1-800-430-5490

ASPIRIN 325 MG
 25 Packets (2 Tablets per Packet)
 Total quantity 50 Tablets

7722-01 Rev C.pdf 1 9/5/2018 2:18:46 PM



Ammonia
Principal Display Panel

796006 Rev. E Unit Carton Printing Plate for "A" size carton.



796006 Rev. E (page 3 of 3)

Principal Display Panel



**Burn Relief Spray
Principal Display Panel**

SHAKE WELL BEFORE USING

Honeywell BURN SPRAY

Water soluble
Benzethonium chloride
Topical antiseptic
Benzocaine
Topical anesthetic
Menthol
Topical anesthetic

Provides antiseptic treatment
and helps relieve the pain of minor burns
and sunburn.

CAUTION: FLAMMABLE
Contents under pressure
Read warning on back panel.

NET WT. 3 OZ (85gm.)

47.0306

Cat. No. 201005

DRUG FACTS

Active ingredients

Benzethonium chloride 0.2% w/w	Topical antiseptic
Benzocaine 1.0% w/w	Topical anesthetic
Menthol 3.3%	Topical anesthetic

Purpose

Uses • for the temporary relief of pain and itching and helps to protect against infection in
• minor cuts and scrapes • burns • sunburn • insect bites • minor skin irritations

Warnings

For external use only

Flammable • keep away from fire or flame • contents under pressure
• do not puncture or incinerate container • do not expose to temperatures above 120°F

Do not use • in or near eyes or other mucus membranes • in case of serious burns
• in case of deep or puncture wounds • for a prolonged period of time
• on large portion of the body

Stop use and ask a doctor if:

• conditions worsens or symptoms persist for more than 7 days
• condition clears up and recurs within a few days
• redness, swelling or irritation occurs

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

• clean the affected area • shake can well before using
• hold 4-6 inches from surface and spray area until wet
• may be covered with a sterile bandage. If bandaged, let dry first
• for adult institutional use only • not intended for use on children

Other information

• avoid inhaling • use only as directed
• intentional misuse by deliberately concentrating and inhaling the contents may be harmful or fatal

Inactive ingredients dipropylene glycol, isobutane, n-butane, propane

Questions or comments? 1-800-430-5490



Distributed by
Honeywell Safety
Products USA, Inc.
Smithfield, RI 02917

Honeywell

4032 Kit Label
340001FP

First Aid

Distributed by Honeywell Safety Products USA, Inc Smithfield, RI 02917

4032 FIRST AID KIT

4032 first aid kit kit

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0498-4032
---------------------	----------------	---------------------------	---------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-4032-01	1 in 1 KIT; Type 0: Not a Combination Product	10/18/2018	

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	1 BOTTLE	30 mL
Part 2	6 PACKET	8.4 mL
Part 3	12 PACKET	24
Part 4	4 PACKET	3.6 g
Part 5	2 AMPULE	0.6 mL
Part 6	1 BOTTLE	59 mL

Part 1 of 6

EYESALINE EMERGENCY EYEWASH

purified water liquid

Product Information

Item Code (Source)	NDC:0498-0100
---------------------------	---------------

Route of Administration	OPHTHALMIC
--------------------------------	------------

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
WATER (UNII: 059QF0KO0R) (WATER - UNII:059QF0KO0R)	WATER	98.6 mL in 100 mL

Inactive Ingredients

Ingredient Name	Strength
SODIUM PHOSPHATE, MONOBASIC, MONOHYDRATE (UNII: 593YOG76RN)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM PHOSPHATE, DIBASIC (UNII: GR686LBA74)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-0100-01	30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part349	12/18/2018	

Part 2 of 6

ANTISEPTIC TOWELETTE

benzalkonium chloride liquid

Product Information

Item Code (Source)	NDC:0498-0501
Route of Administration	TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZALKONIUM CHLORIDE (UNII: F5UM2KM3W7) (BENZALKONIUM - UNII:7N6JUD5X6Y)	BENZALKONIUM CHLORIDE	1.3 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-0501-00	1.4 mL in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part333E	12/21/2017	

Part 3 of 6

ASPIRIN

aspirin tablet

Product Information

Item Code (Source) NDC:0498-0114

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

Inactive Ingredients

Ingredient Name	Strength
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
STARCH, CORN (UNII: O8232NY3SJ)	
POVIDONE (UNII: FZ989GH94E)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
HYPROMELLOSE 2208 (100 MPA.S) (UNII: B1QE5P712K)	
MINERAL OIL (UNII: T5L8T28FGP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	

Product Characteristics

Color	white	Score	2 pieces
Shape	ROUND	Size	10 mm
Flavor		Imprint Code	FR21
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-0114-01	2 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	09/18/2018	

Part 4 of 6

TRIPLE ANTIBIOTIC

bacitracin zinc, polymyxin b sulfate, neomycin sulfate ointment

Product Information

Item Code (Source) NDC:0498-0750

Route of Administration TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BACITRACIN ZINC (UNII: 89Y4M234ES) (BACITRACIN - UNII:58H6RWO52I)	BACITRACIN	400 [iU] in 1 g
POLYMYXIN B SULFATE (UNII: 19371312D4) (POLYMYXIN B - UNII:J2VZ07J96K)	POLYMYXIN B	5000 [iU] in 1 g
NEOMYCIN SULFATE (UNII: 057Y626693) (NEOMYCIN - UNII:I16QD7X297)	NEOMYCIN	3.5 mg in 1 g

Inactive Ingredients

Ingredient Name	Strength
PETROLATUM (UNII: 4T6H12BN9U)	

Product Characteristics

Color	white	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-0750-35	0.9 g in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part333B	09/19/2018	

Part 5 of 6

AMMONIA INHALENT

ammonia inhalent inhalant

Product Information

Item Code (Source) NDC:0498-3334

Route of Administration RESPIRATORY (INHALATION)

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
AMMONIA (UNII: 5138Q19F1X) (AMMONIA - UNII:5138Q19F1X)	AMMONIA	0.045 g in 0.3 mL

Inactive Ingredients

Ingredient Name	Strength
ALCOHOL (UNII: 3K9958V90M)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-3334-00	0.3 mL in 1 AMPULE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		09/18/2018	

Part 6 of 6

BURN RELIEF

lidocaine hydrochloride spray

Product Information

Item Code (Source)	NDC:0498-0221
Route of Administration	TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LIDOCAINE HYDROCHLORIDE (UNII: V13007Z41A) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE HYDROCHLORIDE ANHYDROUS	24.64 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
EDETATE DISODIUM (UNII: 7FLD91C86K)	
TROLAMINE (UNII: 9O3K93S3TK)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
WATER (UNII: 059QF0KO0R)	

TEA TREE OIL (UNII: VIF565UC2G)	
METHYL PARABEN (UNII: A2I8C7HI9T)	
DIAZOLIDINYL UREA (UNII: H5RIZ3MPW4)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
GLYCERIN (UNII: PDC6A3C0OX)	
OCTOXYNOL-9 (UNII: 7JPC6Y25QS)	

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-0221-59	59 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part348	09/18/2018	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		10/18/2018	

Labeler - Honeywell Safety Products USA, INC (079287321)

Establishment			
Name	Address	ID/FEI	Business Operations
James Alexander		040756421	manufacture(0498-3334)

Establishment			
Name	Address	ID/FEI	Business Operations
Honeywell Safety Products USA, INC		079287321	pack(0498-4032)

Establishment			
Name	Address	ID/FEI	Business Operations
Ultra Seal Corporation		085752004	manufacture(0498-0114)

Establishment			
Name	Address	ID/FEI	Business Operations
Water-Jel Technologies		155522589	manufacture(0498-0750, 0498-0221)

Establishment			
Name	Address	ID/FEI	Business Operations
Honeywell Safety Products USA, Inc.		167518617	manufacture(0498-0100)

Establishment

Name	Address	ID/FEI	Business Operations
Changzhou Maokang Medical		421317073	manufacture(0498-0501)

Revised: 3/2019

Honeywell Safety Products USA, INC